Tree Inventory and Preservation Plan Report 5868 County Road 65 Port Hope, Ontario

prepared for

Hillstreet Developments Ltd. 524 Rosebank Road Pickering, ON L1W 2W5

prepared by



146 Lakeshore Road West PO Box 1267 Lakeshore W PO Oakville ON L6K 0B3 t: 289.837.1871 e: consult@kuntzforestry.ca

28 March 2023, revised 7 November 2023 and 2 April 2024

KUNTZ FORESTRY CONSULTING INC Project P3360

Introduction

Kuntz Forestry Consulting Inc. was retained by Hillstreet Developments Ltd. to complete a Tree Inventory and Preservation Plan report as part of a development application for the property located at 5868 County Road 65 in Port Hope. The property is located southwest of County Road 65, within a rural/agricultural area.

The work plan for this tree preservation study included the following:

- Prepare inventory of the individual tree resources over 10cm diameter at breast height (DBH) and trees of all diameters within the road right-of-way on and within six metres of the disturbance limit,
- Evaluate potential tree saving opportunities based on proposed development plans; and
- Document the findings in a Tree Inventory and Preservation Plan Report.

The results of the evaluation are provided below.

Methodology

The tree inventory was conducted on 16 and 24 March 2023. Tree resources were located using the topographic survey and KFCI's Trimble GPS unit, accurate to ~30cm. Individual trees, and trees generally within 6m along the peripheries of the originally expected disturbance areas with the potential to be impacted by the work, were tagged using the numbers 1-255. Individual trees that could not be tagged were identified as Trees A-E. Two Butternut trees were identified as Bn1 and Bn2.

Individual tree resources were assessed utilizing the following parameters:

Tree # - number assigned to tree that corresponds to Figure 1.

Species - common and botanical names provided in the inventory table (Table 1).

DBH - diameter (centimetres) at breast height, measured at 1.4 m above the ground.

Condition - condition of tree considering trunk integrity, crown structure, and crown vigour. Condition ratings include poor (P), fair (F) and good (G).

Crown width – extent of crown (m).

Comments - additional relevant detail. Defects are rated as light (L), moderate (M), or heavy (H).

Polygons (groups of trees, especially forested units) were identified as P1-P13. Descriptions for P1, P9, P12, and P13 can be found within Table 1. Tree polygons P2, P3, P5, P6, P10, and P11 were inventoried by 100% tally, counting all trees within these units and categorizing them by species, size category, and condition [AGS (Acceptable Growing Stock) and UGS (Unacceptable Growing Stock)].

P4, P7, and P8 (larger forested units) were assessed by utilizing fixed area sampling plots (3-4 plots within each unit) and counting all trees within the plots and categorizing them by species, size category, and condition [AGS (Acceptable Growing Stock) and UGS (Unacceptable Growing Stock)].

Tree locations are shown on Figure 1. See Tables 1 and 2 for the results of the inventory. Please note that since the original version of this report, many of these trees are no longer within proximity to the disturbance areas due to changes to the site plan.

Existing Site Conditions

The subject property is currently occupied by agricultural lands, natural heritage features, and a homestead. A larger natural heritage feature exists to the west of the site, and this feature is contiguous with the natural heritage features that exist on-site. Tree resources exist in the form of natural feature trees, individual landscape trees, and hedgerow features. Refer to Figure 1 for the existing conditions.

Tree Resources

The inventory documented 260 individual trees and 13 tree polygons on and within six metres of the subject area. Refer to Tables 1 and 2 for the full tree inventory and Figure 1 for the locations of trees reported in the tree inventory.

Tree resources were comprised of Manitoba Maple (*Acer negundo*), Black Walnut (*Juglans nigra*), Apple species (*Malus sp.*), Trembling Aspen (*Populus tremuloides*), Black Cherry (*Prunus serotina*), Cherry species (*Prunus sp.*), White Pine (*Pinus strobus*), Eastern White Cedar (*Thuja occidentalis*), Sugar Maple (*Acer saccharum*), Silver Maple (*Acer saccharinum*), Red Maple (*Acer rubrum*), Norway Maple (*Acer platanoides*), White Birch (*Betula papyrifera*), White Elm (*Ulmus americana*), Green Ash (*Fraxinus pennsylvanica*), Ironwood (*Ostrya virginiana*), Basswood (*Tilia americana*), American Beech (*Fagus grandifolia*), Red Oak (*Quercus rubra*), Eastern Hemlock (*Tsuga canadensis*), Black Locust (*Robinia pseudoacacia*), Scots Pine (*Pinus sylvestris*), Butternut (*Juglans cinerea*), Yellow Birch (*Betula alleghaniensis*), White Ash (*Fraxinus americana*), Blue Beech (*Carpinus caroliniana*), and Pin Cherry (*Prunus pensylvanica*),

Proposed Development

The proposed development involves the construction of a 40-unit subdivision with single detached dwellings, serviced by central roadways connecting to County Road 65. Grading and servicing, including septic systems for each lot, outlets, and swales will also be required. Refer to Figure 1 for the proposed site plan.

Discussion

The following sections provide a discussion and analysis of development impacts, tree removal requirements, and tree preservation relative to the proposed development and existing conditions.

Development Impacts/Tree Removals

The proposed development will require the removal of 30 trees including, Trees 202-230 and B. These trees conflict directly with the proposed development and related grading.

Refer to Figure 1 for the location of tree removals.

Tree Preservation

The preservation of Trees 1-201, 231-255, A, C-E, BN1, BN2, and P1-P13 will be possible with the use of appropriate tree protection measures as indicated on Figure 1. Tree protection measures will have to be implemented prior to earthworks to ensure designated

tree resources are not impacted by the development. Refer to Figure 1 for the location of required tree preservation fencing and further tree protection plan notes. All grading and disturbances should be directed outside of the TPZ indicated on Figure 1.

A standard tree protection fencing detail is shown on Figure 1 (snow fencing on wooden frame). Alternatively, protection fencing can also be comprised of erosion and sediment control fencing, erected on t-bars and/or affixed paige wire fencing.

<u>Butternut</u>

Two Butternut trees (Bn1 and Bn2) were identified while on site. Pure Butternut (*Juglans cinerea*) are listed as "endangered" per COSEWIC and are protected by the Endangered Species Act (ESA). These trees can be preserved in the context of the development but because work is occurring within 25m, a formal Butternut Health Assessment (BHA) has been conducted. The results will be submitted to the Ministry of Environment, Conservation, and Parks (MECP). Until the BHA is submitted and/or ESA requirements have been satisfied, activity within 25m of these trees is not permitted. Under the ESA, these trees have root harm prevention zone of 9m, and an additional 5m of protection beyond that is required. As these setbacks are being respected (see Figure 1a), additional action under the ESA following the submission of the BHA will not be required.

Summary and Recommendations

Kuntz Forestry Consulting Inc. was retained by Hillstreet Developments Ltd. to complete a Tree Inventory and Preservation Plan report as part of a development application for 5868 County Road 65 in Port Hope. A tree inventory was conducted and reviewed in the context of the proposed development plan.

The findings of the study indicate a total of 260 individual trees and 13 tree polygons on and within six metres of the subject property. The removal of 30 trees is required to accommodate the proposed development. All other tree resources can be saved provided appropriate tree protection measures are installed prior to construction.

The following recommendations are suggested to minimize impacts to trees identified for preservation. Refer to Figure 1 for additional tree preservation notes.

- Tree protection barriers and fencing should be erected at locations prescribed on Figure 1.
- Tree protection measures will have to be implemented prior to construction to ensure the trees identified for preservation are not impacted by the development.
- Branches and roots that extend past prescribed tree protection zones that require pruning must be pruned by a qualified Arborist or other tree professional. All pruning of tree roots and branches must be in accordance with good arboricultural standards.
- Site visits, pre, during and post construction are recommended by either a certified consulting arborist (I.S.A.) or registered professional forester (R.P.F.) to ensure proper utilization of tree protection barriers. Trees should also be inspected for damage

incurred during construction to ensure appropriate pruning or other mitigation measures are implemented.

Respectfully Submitted,

Kuntz Forestry Consulting Inc.

Celine Batterink

Celine Batterink, H.B.Sc. Ecology Senior Consulting Arborist, Ecologist ISA Certified Arborist #ON1546-A Email: <u>cbatterink@kuntzforestry.ca.</u> Phone: 289-837-1871 ext 18

Limitations of Assessment

Only the tree(s) identified in this report were included in the inventory. The assessment of the trees presented in this report has been made using accepted arboricultural techniques. These may include a visual examination taken from the ground of all the above-ground parts of the tree for structural defects, scars, external indications of decay such as fungal fruiting bodies, evidence of attack by insects, discoloured foliage, the condition of any visible root structures, the degree of lean (if any), the general condition of the trees and the identification of potentially hazardous trees or recommendations for removal (if applicable). Where trees could not be directly accessed (ie. due to obstructions, and/or on neighbouring properties), trees were assessed as accurately as possible from nearby vantage points.

Locations of trees provided in the report are determined as accurately as possible based on the best information available. If official survey information is not provided, tree locations in the report may not be exact. Where KFCI's in-house GPS unit is used (if applicable), tree locations are accurate only to the extent that the technology allows, which can be variable based on satellite available, RTK network / cell coverage, canopy coverage, and/or projection transformation limitations. If trees occur on or near property boundaries, an official site survey may be required to determine ownership utilizing specialized survey protocol to gain precise location.

Furthermore, recommendations made in this report are based on the site plans that have been provided at the time of reporting. These recommendations may no longer be applicable should changes be made to the site plan and/or grading, servicing, or landscaping plans following report submission.

Notwithstanding the recommendations and conclusions made in this report, it must be recognized that trees are living organisms, and their health and vigor constantly change over time. They are not immune to changes in site conditions or seasonal variations in the weather conditions. Any tree will fail if the forces applied to the tree exceed the strength of the tree or its parts.

Although every effort has been made to ensure that this assessment is reasonably accurate, the trees should be re-assessed periodically. The assessment presented in this report is valid at the time of inspection.

Table 1. Tree Inventory

Location: 5868 County Rd 65, Port Hope

Date: 16 and 24 March 2023

Surveyors: CB, SA

| Tree# | Common | Scientific Name | DBH | CW | TI | CS | CV | CDB | Comments | Action |
|-------|-----------------|---------------------|-----------|----|-----|-----|-----|-----|---|--------|
| 1 | Trembling Aspen | Populus tremuloides | 22 | 7 | G | G | G | | Deadwood (L) | Retain |
| 2 | Trembling Aspen | Populus tremuloides | 22.5, ~9 | 7 | G | F-G | G | | Deadwood (L) | Retain |
| 3 | Trembling Aspen | Populus tremuloides | 26 | 6 | G | G | G | | Deadwood (L) | Retain |
| 4 | Trembling Aspen | Populus tremuloides | 21 | 6 | G | G | G | | Deadwood (L) | Retain |
| 5 | Sugar Maple | Acer saccharum | 11.5 | 3 | F-G | F-G | G | | Understory tree, V-union at 2m with included bark (L) | Retain |
| 6 | Trembling Aspen | Populus tremuloides | 12 | 3 | G | F | F | | Crooks (M), Deadwood (M) | Retain |
| 7 | Sugar Maple | Acer saccharum | 8.5 | 2 | G | G | G | | | Retain |
| 8 | Sugar Maple | Acer saccharum | 10.5 | 4 | G | G | G | | | Retain |
| 9 | Ironwood | Ostrya virginiana | 11.5 | 5 | G | G | G | | | Retain |
| 10 | White Birch | Betula papyrifera | 9 | 4 | G | G | G | | | Retain |
| 11 | Sugar Maple | Acer saccharum | 12.5, 7.5 | 5 | F | F-G | F | | Union at 0.5, Epicormic branching (L) | Retain |
| 12 | American Beech | Fagus grandifolia | 16.5 | 6 | F | F | P-F | | Beech bark disease (M) | Retain |
| 13 | Trembling Aspen | Populus tremuloides | 36 | 5 | F-G | F | F | 20 | Lean (L), Poor form (M), Asymmetrical crown (M), Deadwood (M) | Retain |
| 14 | Trembling Aspen | Populus tremuloides | 31.5 | 9 | G | F-G | G | | Deadwood (L), Poor form (L) | Retain |
| 15 | Ironwood | Ostrya virginiana | 9 | 4 | G | G | G | | | Retain |
| 16 | Ironwood | Ostrya virginiana | 13 | 3 | F-G | F | F-G | | Asymmetrical crown (M), Lost leader | Retain |
| 17 | Sugar Maple | Acer saccharum | 27 | 8 | G | F-G | G | | Deadwood (L), Broken branches(L) | Retain |
| 18 | American Beech | Fagus grandifolia | 15.5 | 5 | P-F | F | Р | | Asymmetrical crown (H), Beech bark disease (H) | Retain |
| 19 | Trembling Aspen | Populus tremuloides | 26 | 8 | P-F | P-F | P-F | | Canker (L), Bowed crown (H), Lost leader, Deadwood (M) | Retain |
| 20 | Trembling Aspen | Populus tremuloides | 31 | 8 | F | F-G | F-G | | Fruiting bodies (M), Crooks (L), Deadwood (M) | Retain |
| 21 | Ironwood | Ostrya virginiana | 10 | 5 | G | G | G | | | Retain |
| 22 | Trembling Aspen | Populus tremuloides | 32.5 | 8 | F | F | F-G | | Deadwood (M), Asymmetrical crown (M), Crook (M), Fruiting bodies (M) | Retain |
| 23 | Trembling Aspen | Populus tremuloides | 28.5 | 9 | F | F | F-G | | Fruiting bodies (M), Asymmetrical crown (M), Deadwood (M), Crook (M) | Retain |
| 24 | Trembling Aspen | Populus tremuloides | ~32 | 7 | F-G | F-G | F-G | | Fruiting bodies (M), Deadwood (L), Crook (L) | Retain |
| 25 | Sugar Maple | Acer saccharum | ~10 | | G | G | G | | Crowded by 24 | Retain |
| 26 | Trembling Aspen | Populus tremuloides | 25 | 6 | F | F | F | | Fruiting bodies (M), Crook (M), Deadwood (L) | Retain |
| 27 | Trembling Aspen | Populus tremuloides | 28.5 | 7 | F | F-G | F-G | | Fruiting bodies (M), Deadwood (L), Crook (L), Asymmetrical crown (L) | Retain |

| 28 | Trembling Aspen | Populus tremuloides | 29.5 | 6 | F | F | F | Seam with rot (M), Asymmetrical crown (M), Deadwood (M), Fruiting bodies (M) | Retain |
|----|-----------------|---------------------|------|----|-----|-----|-----|---|--------|
| 29 | Trembling Aspen | Populus tremuloides | 31 | 7 | F | G | G | Fruiting bodies (M), Deadwood (M) | Retain |
| 30 | Trembling Aspen | Populus tremuloides | 27 | 7 | F-G | F-G | G | Crook (L), Deadwood (L), Fruiting bodies (L) | Retain |
| 31 | Sugar Maple | Acer saccharum | 22 | 8 | G | G | G | | Retain |
| 32 | Black Cherry | Prunus serotina | 10 | 4 | F | F | F | Bowed (M) over subject property | Retain |
| 33 | Trembling Aspen | Populus tremuloides | 22.5 | 5 | F | P-F | F | Asymmetrical crown (H), Fruiting bodies (L), Cavity (L), Lost leader, Deadwood (L) | Retain |
| 34 | Trembling Aspen | Populus tremuloides | 16.5 | 5 | F | P-F | F | Crook (H), Deadwood (L), Asymmetrical crown (H), stem wound with burl (H) | Retain |
| 35 | Trembling Aspen | Populus tremuloides | 20.5 | 7 | G | G | G | Deadwood (L) | Retain |
| 36 | Trembling Aspen | Populus tremuloides | 13.5 | 5 | G | F-G | F-G | Lean (L), Bowed crown (L) | Retain |
| 37 | Sugar Maple | Acer saccharum | 10 | 5 | G | G | G | | Retain |
| 38 | Sugar Maple | Acer saccharum | 23 | 8 | F-G | F-G | G | V-union in crown, Epicormic branching (L) | Retain |
| 39 | Sugar Maple | Acer saccharum | 17 | 8 | G | G | G | | Retain |
| 40 | Sugar Maple | Acer saccharum | 13.5 | 4 | F-G | F-G | G | Poor form (M), PU in crown | Retain |
| 41 | Sugar Maple | Acer saccharum | 15.5 | 8 | G | G | G | | Retain |
| 42 | White Pine | Pinus strobus | 34 | 12 | F | F | G | Asymmetrical crown (L), co in crown | Retain |
| 43 | Sugar Maple | Acer saccharum | 12.5 | 4 | G | F-G | G | Asymmetrical crown (L) | Retain |
| 44 | Sugar Maple | Acer saccharum | 9.5 | 3 | G | G | G | | Retain |
| 45 | Black Cherry | Prunus serotina | 18 | 7 | G | F-G | G | Asymmetrical crown (L) | Retain |
| 46 | Trembling Aspen | Populus tremuloides | 47 | 14 | P-F | F | F | Fruiting bodies (H), Crooks (H), Deadwood (M) | Retain |
| 47 | Sugar Maple | Acer saccharum | 13 | 4 | G | G | G | | Retain |
| 48 | Sugar Maple | Acer saccharum | 13 | 6 | G | F-G | G | Asymmetrical crown (L) | Retain |
| 49 | Sugar Maple | Acer saccharum | 23 | 6 | F-G | F-G | G | Asymmetrical crown (L), Included fence M) | Retain |
| 50 | Trembling Aspen | Populus tremuloides | 41.5 | 14 | G | G | G | Deadwood (L) | Retain |
| 51 | Trembling Aspen | Populus tremuloides | 16 | 7 | F | F | F-G | Bowed (M), Deadwood (L) | Retain |
| 52 | Sugar Maple | Acer saccharum | 16 | 5 | G | F-G | G | Asymmetrical crown (M) | Retain |
| 53 | Sugar Maple | Acer saccharum | 11 | 5 | G | F-G | G | Asymmetrical crown (L) | Retain |
| 54 | Sugar Maple | Acer saccharum | 10 | 4 | G | G | G | | Retain |
| 55 | Ironwood | Ostrya virginiana | 11 | 7 | G | G | G | | Retain |
| 56 | Sugar Maple | Acer saccharum | 8 | 3 | G | G | G | | Retain |
| 57 | Sugar Maple | Acer saccharum | 12 | 4 | G | G | G | | Retain |
| 58 | Sugar Maple | Acer saccharum | 26 | 12 | G | F-G | G | Asymmetrical crown (L) | Retain |
| 59 | Sugar Maple | Acer saccharum | 79.5 | 14 | P-F | P-F | P-F | Basal rot (H), white rot, Deadwood (L), Lean (L), Asymmetrical crown (M) | Retain |
| 60 | Sugar Maple | Acer saccharum | 13 | 6 | G | G | G | | Retain |

| 61 | Ironwood | Octavo virginiono | 9 | 4 | G | G | G | 1 | | Datain |
|----|-----------------|---------------------|--------------------|----|-----|-----|-----|----|--|--------|
| - | Ironwood | Ostrya virginiana | | | F-G | F | F-G | | | Retain |
| 62 | Basswood | Tilia americana | 21 | 6 | - | - | - | | Crooks (M), Poor form (M) | Retain |
| 63 | Sugar Maple | Acer saccharum | 12 | 3 | G | F-G | G | | Asymmetrical crown (M) | Retain |
| 64 | Sugar Maple | Acer saccharum | 31 | 8 | G | F-G | G | | Asymmetrical crown (M) | Retain |
| 65 | Trembling Aspen | Populus tremuloides | 18.5 | 5 | P-F | P-F | P-F | 35 | Fruiting bodies (M), Bowed (H), Deadwood (M) | Retain |
| 66 | White Birch | Betula papyrifera | 33 | 5 | Р | Р | Р | 80 | Deadwood (H), rot (H) | Retain |
| 67 | Trembling Aspen | Populus tremuloides | 32.5 | 8 | F | F | F | | Fruiting bodies (M), Crooks (M), Asymmetrical crown (M), Poor form (M) | Retain |
| 68 | Sugar Maple | Acer saccharum | 23 | 6 | G | F-G | G | | Asymmetrical crown (L) | Retain |
| 69 | Sugar Maple | Acer saccharum | 17 | 8 | G | G | G | | | Retain |
| 70 | Sugar Maple | Acer saccharum | 13 | 5 | F | F-G | G | | Asymmetrical crown (L), stem wounds (M) | Retain |
| 71 | Sugar Maple | Acer saccharum | 11.5 | 4 | G | G | G | | | Retain |
| 72 | Sugar Maple | Acer saccharum | 21.5 | 7 | G | F-G | G | | Asymmetrical crown (L) | Retain |
| 73 | Trembling Aspen | Populus tremuloides | 30.5 | 8 | F | F | F | | Fruiting bodies (M), Crooks (M), Deadwood (L) | Retain |
| 74 | Sugar Maple | Acer saccharum | 14.5 | 6 | G | G | G | | | Retain |
| 75 | Sugar Maple | Acer saccharum | 60 | 12 | F | F | F | | Union at 1m, Poor form (M) Asymmetrical crown (L), Deadwood (L) | Retain |
| 76 | Red Oak | Quercus rubra | ~78, 65, 65, 34 | 30 | F-G | F | F-G | | Union at 0.3m, Asymmetrical crown (M), Deadwood (M), large spreading leaders | Retain |
| 77 | Sugar Maple | Acer saccharum | 27 | 7 | F | G | F-G | | Sugar Maple borer (M) | Retain |
| 78 | Sugar Maple | Acer saccharum | 35 | 7 | G | F | G | | Asymmetrical crown (H) | Retain |
| 79 | Sugar Maple | Acer saccharum | ~39, 38 | 12 | F | F-G | F-G | | V-union at 1m | Retain |
| 80 | Sugar Maple | Acer saccharum | 37.5 | 8 | G | G | G | | | Retain |
| 81 | Sugar Maple | Acer saccharum | 24.5 | 6 | G | F | G | | Asymmetrical crown (M) | Retain |
| 82 | Sugar Maple | Acer saccharum | 40 | 6 | G | F-G | G | | Asymmetrical crown (L) | Retain |
| 83 | Sugar Maple | Acer saccharum | 27.5 | 8 | G | F-G | G | | Asymmetrical crown (L) | Retain |
| 84 | Red Maple | Acer rubrum | 36 | 8 | F-G | F-G | F-G | | 1 dead stem at base, Asymmetrical crown (L) | Retain |
| 85 | American Beech | Fagus grandifolia | 23 | 7 | P-F | G | P-F | | Beech bark disease (H) | Retain |
| 86 | Red Maple | Acer rubrum | 39, 48.5 | 12 | F-G | F-G | G | | Union at 0.2m, Asymmetrical crown (L) | Retain |
| 87 | Sugar Maple | Acer saccharum | 15.5 | 7 | G | G | G | | | Retain |
| 88 | Red Maple | Acer rubrum | 61 | 12 | F | F-G | G | | V-union at 1.6m with included bark (L), Asymmetrical crown (L) | Retain |
| 89 | American Beech | Fagus grandifolia | 20 | 6 | F | G | P-F | 1 | Beech bark disease (M) | Retain |
| 90 | American Beech | Fagus grandifolia | 18, 13 | 8 | F | F | P-F | | V-union at 1m, Beech bark disease (M) | Retain |
| 91 | Black Cherry | Prunus serotina | 40.5 | 7 | G | G | G | | Deadwood (L) | Retain |
| 92 | Ironwood | Ostrya virginiana | 19, 15.5 | 7 | F-G | F | F-G | | Union at base, Asymmetrical crown (M) | Retain |
| 93 | Ironwood | Ostrya virginiana | 15.5, 12 | 7 | F-G | F-G | F-G | İ | Union at 0.2m, Poor form (L), Deadwood (M) | Retain |

| 94 | White Pine | Pinus strobus | 58 | 11 | G | F-G | G | | Deadwood (L), Asymmetrical crown (M) | Retain |
|-----|-----------------|------------------------|---------------|----|-----|-----|-----|-------|---|--------|
| 95 | Ironwood | Ostrya virginiana | 10 | 3 | G | F-G | F-G | | Poor form (M) | Retain |
| 96 | White Pine | Pinus strobus | 53 | 13 | G | G | G | ł – – | | Retain |
| 97 | Sugar Maple | Acer saccharum | 13 | 4 | G | F-G | G | | Lean (L), Asymmetrical crown (L), Crook (L) | Retain |
| 98 | Sugar Maple | Acer saccharum | 47 | 12 | G | F-G | G | | Lean (L), Asymmetrical crown (L) | Retain |
| 99 | Sugar Maple | Acer saccharum | 9.5, 8 | 4 | G | G | G | | Union at 1m | Retain |
| 100 | White Pine | Pinus strobus | 34 | 7 | G | F-G | G | | Asymmetrical crown (L), Deadwood (L) | Retain |
| 101 | Red Oak | Quercus rubra | 57 | 13 | G | F-G | G | | Lean (L), Asymmetrical crown (M) | Retain |
| 102 | Ironwood | Ostrya virginiana | 16 | 3 | F | F | P-F | 70 | Crowded by 101, Asymmetrical crown (H) | Retain |
| 103 | Black Cherry | Prunus serotina | 40 | 12 | G | F-G | G | | Deadwood (L), Asymmetrical crown (L) | Retain |
| 104 | Black Cherry | Prunus serotina | 60 | 18 | G | F | F-G | | Deadwood (L), Epicormic branching (L), Asymmetrical crown (L) | Retain |
| 105 | Sugar Maple | Acer saccharum | 14 | 3 | G | G | G | | | Retain |
| 106 | Green Ash | Fraxinus pennsylvanica | 12.5 | 3 | F | F | F | | Crook (M) | Retain |
| 107 | Sugar Maple | Acer saccharum | 20.5 | 7 | G | G | G | | Deadwood (L) | Retain |
| 108 | Sugar Maple | Acer saccharum | 17.5 | 5 | G | G | G | | Deadwood (L) | Retain |
| 109 | Sugar Maple | Acer saccharum | 20.5 | 8 | G | G | G | | | Retain |
| 111 | Sugar Maple | Acer saccharum | 18 | 6 | F | G | G | | Stem wound from rubbing against 112 (M) | Retain |
| 112 | Trembling Aspen | Populus tremuloides | 28 | 7 | F | F | F | | Lean (M), Crooks (M), Deadwood (M) | Retain |
| 113 | Sugar Maple | Acer saccharum | 11 | 3 | G | G | G | | | Retain |
| 114 | Sugar Maple | Acer saccharum | 18.5 | 8 | G | G | G | | | Retain |
| 115 | Trembling Aspen | Populus tremuloides | 31 | 4 | F-G | F | F | | Asymmetrical crown (M), Deadwood (M), Crooks (L) | Retain |
| 116 | Black Cherry | Prunus serotina | 20 | 5 | F-G | F | F-G | | Bowed (M) | Retain |
| 117 | Sugar Maple | Acer saccharum | 24 | 9 | G | F-G | G | | Asymmetrical crown (L) | Retain |
| 118 | Sugar Maple | Acer saccharum | 27 | 7 | G | F-G | G | | Asymmetrical crown (L) | Retain |
| 119 | Trembling Aspen | Populus tremuloides | 24.5 | 8 | G | G | G | | | Retain |
| 120 | Sugar Maple | Acer saccharum | 14.5 | 4 | G | F-G | G | | Crowded by 119 | Retain |
| 121 | Trembling Aspen | Populus tremuloides | 24 | 7 | G | F-G | G | | Asymmetrical crown (L) | Retain |
| 122 | Black Cherry | Prunus serotina | 20.5 | 8 | F-G | F-G | G | | Crook (M) | Retain |
| 123 | Black Cherry | Prunus serotina | 26, 21, 17 | 10 | F | F | F | | Union at 0.5 and 1m, Lost leader, burl, Poor form (M) | Retain |
| 124 | Black Locust | Robinia pseudoacacia | 18 | 7 | G | G | G | | | Retain |
| 125 | Sugar Maple | Acer saccharum | 17, 20 | 10 | F | F-G | G | | V-union at .3m with included bark (L) | Retain |
| 126 | White Birch | Betula papyrifera | 23.5 | 11 | F-G | F-G | G | | Bowed (L), Deadwood (L) | Retain |
| 127 | Sugar Maple | Acer saccharum | 25 | 10 | G | G | G | | | Retain |
| 128 | White Birch | Betula papyrifera | 24 | 10 | F-G | F-G | G | | Stem wounds (L), Deadwood (L), Lean (L) | Retain |
| 129 | White Birch | Betula papyrifera | 28, 24 | 14 | F | F | F-G | | Fruiting bodies (L), Union at base, Deadwood (L) | Retain |
| 130 | Sugar Maple | Acer saccharum | 25 | 8 | G | G | G | | | Retain |

| 131 | Silver Maple | Acer saccharinum | 22.5 | 5 | G | G | G | | | Retain |
|-----|-----------------|-------------------|----------|----|-----|-----|-----|----|--|--------|
| 132 | Basswood | Tilia americana | 18.5 | 6 | G | F | F-G | | Grapevine competition (M), Bowed crown (H) | Retain |
| 133 | Eastern Hemlock | Tsuga canadensis | ~32 | 8 | G | G | G | | | Retain |
| 134 | White Birch | Betula papyrifera | 23 | 8 | G | G | G | | | Retain |
| 135 | White Birch | Betula papyrifera | 16.5 | 4 | G | F | F-G | | Poor form (L) | Retain |
| 136 | Cherry species | Prunus sp. | 21 | 6 | F | F | F | 40 | Deadwood (M) | Retain |
| 137 | White Birch | Betula papyrifera | 21 | 7 | F | F | F | | Lean (M), Deadwood (L) | Retain |
| 138 | Ironwood | Ostrya virginiana | 11.5 | 5 | G | F-G | G | | Asymmetrical crown (L) | Retain |
| 139 | Red Maple | Acer rubrum | 17 | 6 | G | G | G | | | Retain |
| 140 | White Birch | Betula papyrifera | 31 | 10 | F-G | F-G | F-G | | Deadwood (L), Bowed (L), Asymmetrical crown (L) | Retain |
| 141 | White Birch | Betula papyrifera | 14 | 5 | G | F-G | G | | Bowed (L) | Retain |
| 142 | White Birch | Betula papyrifera | 30 | 9 | G | G | G | | Deadwood (M), Asymmetrical crown (L) | Retain |
| 143 | White Birch | Betula papyrifera | 29.5 | 8 | G | G | G | | Deadwood (L) | Retain |
| 144 | American Beech | Fagus grandifolia | 11 | 3 | F | F | P-F | | Beech bark disease (M) | Retain |
| 145 | White Birch | Betula papyrifera | 31, 15 | 10 | F-G | F-G | F-G | | Union at 0.2m, Deadwood (L) | Retain |
| 146 | Sugar Maple | Acer saccharum | 13 | 6 | G | G | G | | | Retain |
| 147 | Red Oak | Quercus rubra | 12.5 | 6 | G | G | G | | | Retain |
| 148 | Red Maple | Acer rubrum | 32.5 | 8 | G | G | G | | | Retain |
| 149 | Red Maple | Acer rubrum | 12.5 | 4 | G | F-G | G | | Asymmetrical crown (M) | Retain |
| 150 | Apple sp | Malus sp | 40 | 6 | P-F | Р | Р | 70 | Deadwood (H), Epicormic branching (M), Bowed (M) | Retain |
| 151 | Apple sp. | Malus sp | 29 | 7 | P-F | P-F | P-F | 50 | Deadwood (H), Epicormic branching (M) | Retain |
| 152 | Sugar Maple | Acer saccharum | 18 | 6 | G | G | G | | | Retain |
| 153 | Black Cherry | Prunus serotina | 15.5 | 4 | G | F-G | G | | Deadwood (L), Asymmetrical crown (L) | Retain |
| 154 | White Birch | Betula papyrifera | 29 | 6 | F | F | F-G | | Bowed (M) over subject property, Deadwood (L) | Retain |
| 155 | White Birch | Betula papyrifera | 24 | 4 | Р | F | P-F | | Lean (L), canker (H) | Retain |
| 156 | Black Cherry | Prunus serotina | 24.5 | 7 | F-G | F-G | F-G | | Black knot (L), Asymmetrical crown (L), Deadwood (L) | Retain |
| 157 | tag not used | | | | | | | | | |
| 158 | Silver Maple | Acer saccharinum | 27 | 7 | G | G | G | | | Retain |
| 159 | Red Maple | Acer rubrum | 14 | 5 | G | F-G | G | | Asymmetrical crown (L) | Retain |
| 160 | Sugar Maple | Acer saccharum | 12 | 5 | G | G | G | | | Retain |
| 161 | Apple sp. | Malus sp | ~8, 9, 7 | 7 | P-F | Ρ | P-F | | Union at 1m, Bowed (H) over subject property, Vine competition (H) | Retain |
| 162 | White Birch | Betula papyrifera | 23, 25 | 7 | F-G | F | F-G | | Union at 1.3m, Asymmetrical crown (M), Grapevine competition (L) | Retain |
| 163 | White Birch | Betula papyrifera | 28 | 6 | F-G | G | G | | Crook (L), Bowed (L) | Retain |
| 164 | Sugar Maple | Acer saccharum | 17 | 5 | G | G | G | | | Retain |

| 165 | White Birch | Betula papyrifera | 26, 25 | 8 | F | F-G | F | | V-union at 0.4m with included bark (L) and stem wound (M), Deadwood (L), Asymmetrical crown (L) | Retain |
|-----|---------------------|----------------------|----------------|-----|-----|-----|-----|----|--|--------|
| 166 | Red Oak | Quercus rubra | 38, 22.5 | 9 | F | F-G | F-G | | Lean (L), Union at 0.3m, Asymmetrical crown (L) | Retain |
| 167 | Red Maple | Acer rubrum | 28.5 | 8 | G | F-G | F-G | | Asymmetrical crown (L), Grapevine competition (L), Poor form (L) | Retain |
| 168 | Black Cherry | Prunus serotina | 10.5 | 2 | F | F | F | | Crooks (M), Epicormic branching (L), Poor form (L) | Retain |
| 169 | Sugar Maple | Acer saccharum | 30.5 | 8 | F | F-G | F-G | | V-union at 3m with included bark (L) | Retain |
| 170 | Red Oak | Quercus rubra | 30.5 | 10 | F-G | F-G | G | | Lean (L), stem wound (L), Asymmetrical crown (L) | Retain |
| 171 | Sugar Maple | Acer saccharum | 20 | 3 | G | G | G | | | Retain |
| 172 | White Birch | Betula papyrifera | 31, ~23, 22 | 11 | F | F | F | 20 | 1 dead stem, Union at base, Deadwood (M) | Retain |
| 173 | American Beech | Fagus grandifolia | 19.5 | 6 | G | G | G | | Asymmetrical crown (L) | Retain |
| 174 | Sugar Maple | Acer saccharum | 10.5 | 4 | G | G | G | | | Retain |
| 175 | Sugar Maple | Acer saccharum | 22 | 8 | G | G | G | | | Retain |
| 176 | White Pine | Pinus strobus | 24 | 6 | G | G | G | | | Retain |
| 177 | Sugar Maple | Acer saccharum | 18 | 5 | G | G | G | | | Retain |
| 178 | White Birch | Betula papyrifera | 12.5 | 4 | G | G | G | | Lean (L) | Retain |
| 179 | White Birch | Betula papyrifera | 13.5 | 5 | G | F-G | G | | Asymmetrical crown (L) | Retain |
| 180 | White Birch | Betula papyrifera | 10 | 3 | G | G | G | | | Retain |
| 181 | White Birch | Betula papyrifera | 10.5 | 3 | G | G | G | | | Retain |
| 182 | White Birch | Betula papyrifera | 14.5 | 4 | G | G | G | | Lean (L) | Retain |
| 183 | Black Locust | Robinia pseudoacacia | 24 | 10 | G | F-G | G | | Asymmetrical crown (L) | Retain |
| 184 | Black Locust | Robinia pseudoacacia | 19 | 10 | G | G | G | | Broken branches(L) | Retain |
| 185 | Black Locust | Robinia pseudoacacia | 20.5 | 10 | G | G | G | | | Retain |
| 186 | Black Locust | Robinia pseudoacacia | ~18 | 10 | F-G | F-G | F-G | | Stem wound (L), Grapevine competition (L) | Retain |
| 187 | Black Locust | Robinia pseudoacacia | 12 | 8 | G | F-G | F-G | | Asymmetrical crown (M), Lean (L) | Retain |
| 188 | Black Walnut | Juglans nigra | 11.5 | 4 | G | G | G | | | Retain |
| 189 | Sugar Maple | Acer saccharum | 37 | 12 | G | G | G | | | Retain |
| 190 | Black Locust | Robinia pseudoacacia | 58.5 | 12 | F | F-G | F | | De (L), Fruiting bodies (L) | Retain |
| 191 | Sugar Maple | Acer saccharum | 37, 25.5 | 12 | F | F-G | F-G | | V-union at .8m with included bark (M), Deadwood (L), Grapevine competition (L), Poor form (L) | Retain |
| 192 | Eastern White Cedar | Thuja occidentalis | ~12, 17 | 4 | F | F | F | | Union at base with 1 dead stem | Retain |
| 193 | Eastern White Cedar | Thuja occidentalis | 21 | 4 | F-G | F-G | F-G | | Asymmetrical crown (L), Lean (L) | Retain |
| 194 | Eastern White Cedar | Thuja occidentalis | 11 | 1.5 | G | F-G | G | | Asymmetrical crown (L) | Retain |
| 195 | Eastern White Cedar | Thuja occidentalis | 20 | 4 | F | F | F | | Lean (H) over creek | Retain |
| 196 | Eastern White Cedar | Thuja occidentalis | 25, 13 | 4 | G | F-G | F-G | | Union at 0.3m, Deadwood (L) | Retain |
| 197 | Eastern White Cedar | Thuja occidentalis | 12 | 4 | F-G | G | F-G | | Lean (L) over creek | Retain |

| 198 | Eastern White Cedar | Thuja occidentalis | 10 | 2 | G | G | G | | Retain |
|-----|---------------------|----------------------|-------------------|----|-----|-----|-----|---|--------|
| 199 | Eastern White Cedar | Thuja occidentalis | 18.5, 188 | 4 | G | G | G | Union at base | Retain |
| 200 | Eastern White Cedar | Thuja occidentalis | 15.5 | 3 | F | F | F-G | Sweep (H), Lost leader | Retain |
| 201 | Eastern White Cedar | Thuja occidentalis | ~14 | 5 | G | G | G | | Retain |
| 202 | Sugar Maple | Acer saccharum | 120.5 | 24 | P-F | F-G | F | Fruiting bodies, Union at 2m, rot, Deadwood (M), prune if saving | Remove |
| 203 | Sugar Maple | Acer saccharum | 41 | 10 | F | F-G | G | V-union at 3m with included bark (M), Asymmetrical crown (L) | Remove |
| 204 | Red Maple | Acer rubrum | 36 | 8 | G | F-G | G | Asymmetrical crown (M), Deadwood (L) | Remove |
| 205 | Sugar Maple | Acer saccharum | 21.5 | 9 | G | F-G | G | Asymmetrical crown (M) | Remove |
| 206 | Sugar Maple | Acer saccharum | 33 | 6 | G | F-G | G | Asymmetrical crown (M) | Remove |
| 207 | Sugar Maple | Acer saccharum | 78.5 | 16 | F | F-G | F-G | V-union at 2m | Remove |
| 208 | Manitoba Maple | Acer negundo | 15.5 | 6 | G | G | G | | Remove |
| 209 | Sugar Maple | Acer saccharum | 57 | 12 | F | F-G | G | V-union a 1m with included bark (L) | Remove |
| 210 | Sugar Maple | Acer saccharum | 18 | 6 | G | G | G | | Remove |
| 211 | Sugar Maple | Acer saccharum | 54.5 | 12 | G | G | G | | Remove |
| 212 | Sugar Maple | Acer saccharum | 93 | 20 | F | F-G | F-G | Poor union at 2m, Deadwood (M) | Remove |
| 213 | Red Maple | Acer rubrum | 60.5 | 12 | F | F | F-G | Seams (M), Deadwood (L), Asymmetrical crown (L) | Remove |
| 214 | Silver Maple | Acer saccharinum | 79 | 15 | F | F-G | F | Poor union at 2m with rot, Asymmetrical crown (L) | Remove |
| 215 | Sugar Maple | Acer saccharum | 14.5 | 7 | G | G | G | | Remove |
| 216 | Black Locust | Robinia pseudoacacia | 132 | 20 | Р | P-F | P-F | 1 leader failed at 1m, Deadwood (H), girdling wound (H), FB, rot (H) | Remove |
| 217 | Manitoba Maple | Acer negundo | 17, 24 | 10 | F-G | F-G | G | Union at 0.2m, Coppice Growth (L), Epicormic branching (L) | Remove |
| 218 | Apple sp | Malus sp | ~21, 19, 17 | 7 | F | F | F | Union at 0.5m, Epicormic branching (M), Pruning wounds(L) | Remove |
| 219 | Apple sp | Malus sp | 23.5, 32.5 | 10 | P-F | F | F | Rot (H), Union at 1m, Epicormic branching (M) | Remove |
| 220 | Apple sp | Malus sp | 16, ~21 | 7 | F-G | F-G | F | Union at base, Grapevine competition (L) | Remove |
| 221 | Apple sp | Malus sp | 21.5, ~23, 14 | 8 | P-F | F | F | Union at 1m, hollow, Epicormic branching (L) | Remove |
| 222 | Apple sp | Malus sp | 22, 29 | 7 | P-F | F | F | Union at 0.5m with rot, Epicormic branching (M) | Remove |
| 223 | Apple sp | Malus sp | 14, ~13, 12 | 7 | F | F | P-F | Union at 1m with rot, Sapsucker damage (M), Epicormic branching (L) | Remove |
| 224 | Apple sp | Malus sp | 27, 25.5, 19.5 | 8 | P-F | F | P-F | Union at 0.4m with rot, Epicormic branching (M), Deadwood (M) | Remove |
| 225 | Apple sp | Malus sp | 65.5 | 6 | P-F | P-F | P-F | Union at 1.2m w pruned leader, Poor form (H), Epicormic branching (M) | Remove |

| 226 | Apple sp | Malus sp | 38.5 | 8 | P-F | F | P-F | | Stem wound (H) with rot from failed leader, Epicormic branching (H), Asymmetrical crown (M) | Remove |
|-----|----------------|----------------------|---------------------|----|-----|-----|-----|----|---|--------|
| 227 | Apple sp | Malus sp | 53 | 6 | P-F | F | P-F | | Rot (M), Pruning wounds(H), Epicormic branching (H), Poor form (M) | Remove |
| 228 | Black Locust | Robinia pseudoacacia | 38.5 | 8 | F | G | G | | V-union at 1.2m, Epicormic branching (L) | Remove |
| 229 | Sugar Maple | Acer saccharum | 73.5 | 10 | F | F-G | G | | V-union at 2m, seam (M), Poor form (L) | Remove |
| 230 | Norway Maple | Acer platanoides | 24 | 6 | Р | G | F | | Canker (H) | Remove |
| 231 | Manitoba Maple | Acer negundo | ~65 | 15 | F | P-F | P-F | | Bowed (H), Epicormic branching (H) | Retain |
| 232 | Manitoba Maple | Acer negundo | ~75, 75 | 18 | F | F | F | | Union at 1m, Epicormic branching (H), Bowed (M) | Retain |
| 233 | Manitoba Maple | Acer negundo | 28.5 | 7 | F | F | F-G | | Growing from old stump, Lean (M), Epicormic branching (L) | Retain |
| 234 | Manitoba Maple | Acer negundo | 33.5 | 7 | F | F | F | | Bowed (M) north, Epicormic branching (H), Poor form (H), Broken branches(M) | Retain |
| 235 | Manitoba Maple | Acer negundo | 26 | 6 | F | F | F-G | | Bowed (M) southwest, Epicormic branching (M) | Retain |
| 236 | Manitoba Maple | Acer negundo | 23 | 8 | F-G | F-G | F-G | | Bowed (M) south, Union at 1.7m, Epicormic branching (L) | Retain |
| 237 | Manitoba Maple | Acer negundo | 23, 23, 26, 21.5 | 10 | F | F-G | F-G | | Poor union at base and.5m, ab (M), Bowed (L) | Retain |
| 238 | Manitoba Maple | Acer negundo | 19 | 6 | F | F | F-G | | Bowed (M) southeast, Poor form (M), Epicormic branching (L) | Retain |
| 239 | Sugar Maple | Acer saccharum | 15.5, ~7 | 4 | G | G | G | | Union at .2m | Retain |
| 240 | Manitoba Maple | Acer negundo | 23 | 8 | F | F | F-G | | Bowed (H) south, Epicormic branching (M) | Retain |
| 241 | Black Locust | Robinia pseudoacacia | 109 | 15 | F | F | P-F | 30 | V-union at 1m, Deadwood (H), Asymmetrical crown (M) | Retain |
| 242 | Manitoba Maple | Acer negundo | 27.5, ~45 | 16 | F | P-F | F | | Union at 0.2m and 1m, Bowed (H) south, Epicormic branching (M), Poor form (M), Broken branches(M) | Retain |
| 243 | Black Walnut | Juglans nigra | 11 | 4 | G | G | G | | | Retain |
| 244 | Black Locust | Robinia pseudoacacia | 31 | 7 | F-G | F-G | F-G | | V-union at 1.3m | Retain |
| 245 | Black Locust | Robinia pseudoacacia | ~8, 5 | 4 | G | G | G | | Union at 2m | Retain |
| 246 | Black Locust | Robinia pseudoacacia | 33 | 7 | F-G | G | G | | V-union at 1.1m | Retain |
| 247 | Black Locust | Robinia pseudoacacia | 12 | 5 | G | G | G | | | Retain |
| 248 | Manitoba Maple | Acer negundo | 15, 17, ~15, 15 | 6 | F-G | G | F-G | | Union at.2m, Epicormic branching (M) | Retain |
| 249 | Manitoba Maple | Acer negundo | 20, ~20 | 8 | F | F | F-G | 1 | Union at base, Epicormic branching (L), Poor form (L) | Retain |
| 250 | Manitoba Maple | Acer negundo | 25, ~21 | 7 | F-G | F | F | | Union at base, Epicormic branching (L), Poor form (M), stem wound from branch from Tree D | Retain |
| 251 | Black Walnut | Juglans nigra | 10.5 | 4 | G | G | G | 1 | | Retain |
| 252 | Black Locust | Robinia pseudoacacia | 14 | 3 | G | G | G | | | Retain |
| 253 | Red Maple | Acer rubrum | 10 | 4 | G | G | G | | | Retain |
| 254 | Red Maple | Acer rubrum | 14 | 3 | G | G | G | | | Retain |

| 255 | Basswood | Tilia americana | 25 | 5 | G | F-G | G | | Asymmetrical crown (L) | Retain |
|-----|---------------------------------|---|-------------------|-----|-----|-----|-----|----|--|--------|
| А | Silver Maple | Acer saccharinum | 93.5 | 18 | G | G | G | | | Retain |
| В | Apple sp | Malus sp | 34 | 6 | F | F | F | | Pruning wounds(M), Epicormic branching (M), Deadwood (M), Poor form (M) | Remove |
| С | White Birch | Betula papyrifera | ~17, 16, 14, 9 | 5 | P-F | P-F | P-F | 30 | Deadwood (M), Lost leader's, seam (M), Union at base | Retain |
| D | Scots Pine | Pinus sylvestris | ~45 | 8 | F-G | G | G | | Crook (L) | Retain |
| Е | Black Walnut | Juglans nigra | ~62 | 15 | F-G | F-G | F-G | | V-union at 2m, Deadwood (L) | Retain |
| Bn1 | Butternut | Juglans cinerea | 8 | 5 | F-G | G | G | | Deer rub damage (M) | Retain |
| Bn2 | Butternut | Juglans cinerea | 6.5 | 4 | G | G | G | | | Retain |
| P1 | White Birch, Trembling Aspen | Betula papyrifera, Populus Tremuloides | ~2-9 | 2.0 | G | G | G | | Cluster of approximately 200 White Birch, 80 Trembling Aspen | Retain |
| P2 | See Table 2 | | | | | | | | | Retain |
| P3 | See Table 2 | | | | | | | | | Retain |
| P4 | See Table 2 | | | | | | | | | Retain |
| P5 | See Table 2 | | | | | | | | | Retain |
| P6 | See Table 2 | | | | | | | | | Retain |
| P7 | See Table 2 | | | | | | | | | Retain |
| P8 | See Table 2 | | | | | | | | | Retain |
| P9 | White Birch | Betula papyrifera | <10 | 2.0 | G | G | G | | Pocket of dense regeneration | Retain |
| P10 | See Table 2 | • | • | | • | • | | • | | Retain |
| P11 | See Table 2 | | | | | | | | | Retain |
| P12 | Black Locust | Robinia pseudoacacia | <10 | 2.0 | G | G | G | | 19 trees, regeneration | Retain |
| P13 | Black Locust | Robinia pseudoacacia | ~10-18 | 4.0 | G | G | G | | 20 trees, 3 shared with the right-of-way, also regeneration-sized Black Locust and Sumac within unit | Retain |

| 28 March 2023, revised 7 November 2023, 2 April 2024 |
|--|
|--|

| Codes | | | | | | | | |
|---|---------------------------|-----------|--|--|--|--|--|--|
| DBH | Diameter at Breast Height | (cm) | | | | | | |
| ТІ | Trunk Integrity | (G, F, P) | | | | | | |
| CS | Crown Structure | (G, F, P) | | | | | | |
| CV | Crown Vigor | (G, F, P) | | | | | | |
| CDB | Crown Die Back | (%) | | | | | | |
| CW | Crown Width | (m) | | | | | | |
| ~ = estimate; (VL) = very light; (L) = light; (M) = moderate; (H) = heavy; (VH) = very heavy | | | | | | | | |

Table 2. 100% Tally or Fixed Area Sampling of Polygons

| Location: | |
|-------------------|--|
| Date: | |
| Surveyor: | |
| Compartment: | |
| Stations Tallied: | |

5868 County Road 65 24-Mar-23 SA P2 100% Tally

| Tree Size Class >>>> | | Polewood 10-24 cm | | | | | | | | | |
|-----------------------------|--------------|----------------------|-----|-------------------|-----|--------------------|-----|------------------|-----|-----------------|-----|
| | | | | Small 26-36 cm | | Medium 38-48 cm | | Large 50 cm + | | Total All Sizes | |
| Species | | AGS | UGS | AGS | UGS | AGS | UGS | AGS | UGS | AGS | UGS |
| White Birch (Betula papyrif | era) | 5 | 2 | | | | | | | 5 | 2 |
| Trembling Aspen (Populus | tremuloides) | 2 | 2 | | 2 | | | | | 2 | 4 |
| American Beech (Fagus gr | andifolia) | 6 | 1 | | | | | | | 6 | 1 |
| Sugar Maple (Acer sacchai | rum) | 16 | 5 | | | | | | | 16 | 5 |
| Black Cherry (Prunus serot | ina) | | 2 | | | | | | | 0 | 2 |
| Ironwood (Ostrya virginiana |) | 1 | | | | | | | | 1 | 0 |
| Total Number of | of Trees | 30 | 12 | 0 | 2 | 0 | 0 | 0 | 0 | 30 | 14 |

Description

| Location: | 5868 County Road 65 |
|-------------------|---------------------|
| Date: | 24-Mar-23 |
| Surveyor: | SA |
| Compartment: | P3 |
| Stations Tallied: | 100% Tally |

| | | Delevered | | | | | | | | | |
|-------------------------|-----------------|----------------------|-----|-------------------|-----|--------------------|-----|------------------|-----|-----------------|-----|
| Tree Size | Class >>>> | Polewood 10-24 cm | | Small 26-36 cm | | Medium 38-48 cm | | Large 50 cm + | | Total All Sizes | |
| Spe | ecies | AGS | UGS | AGS | UGS | AGS | UGS | AGS | UGS | AGS | UGS |
| Red Oak (Quercus rub | ora) | 5 | 2 | 3 | | | | | | 8 | 2 |
| White Birch (Betula pa | apyrifera) | 12 | | | | | | | | 12 | 0 |
| Yellow Birch (Betula a | llegheniensis) | 1 | | | | | | | | 1 | 0 |
| Ironwood (Ostrya virgir | niana) | 3 | 1 | | | | | | | 3 | 1 |
| Sugar Maple (Acer sad | ccharum) | 1 | | | | | | | | 1 | 0 |
| | | | | | | | | | | 0 | 0 |
| Total Num | ber of Trees | 22 | 3 | 3 | 0 | 0 | 0 | 0 | 0 | 25 | 3 |

Description

Copious White Birch regeneration in unit

| Location: | 5868 County Road 65 |
|-------------------|---------------------|
| Date: | 24-Mar-23 |
| Surveyor: | SA |
| Compartment: | P4 |
| Stations Tallied: | 3 |

3.99m radius fixed area plots

| | | Polewood | | Sawtimber Sizes | | | | | | | | |
|------------------------|-------------------|----------|-----|-----------------|---|-----|-----|-----|-----|-----------------|-----|--|
| Tree Size | Class >>>> | 10-24 cm | | - | Small Mediu 26-36 cm 38-48 cm | | | | • | Total All Sizes | | |
| Spe | cies | AGS | UGS | AGS | UGS | AGS | UGS | AGS | UGS | AGS | UGS | |
| Trembling Aspen (Pop | ulus tremuloides) | 13 | 1 | | | | | | | 13 | 1 | |
| White Birch (Betula pa | pyrifera) | 2 | | | | | | | | 2 | 0 | |
| | | | | | | | | | | 0 | 0 | |
| | | | | | | | | | | 0 | 0 | |
| Total Numb | per of Trees | 15 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 15 | 1 | |

Description:

Dense amounts of regen-sized trees (<10cm DBH) including Trembling Aspen, Green Ash, White Birch, Black Cherry, and Sugar Maple

| Location: | 5868 County Road 65 |
|-------------------|---------------------|
| Date: | 24-Mar-23 |
| Surveyor: | SA |
| Compartment: | P5 |
| Stations Tallied: | 100% Tally |

| | Dala | Polewood 10-24 cm | | Sawtimber Sizes | | | | | | | |
|--------------------------------|------|----------------------|-----|-----------------|-----|-----|-----|---------------------|-----|-----------|--|
| Tree Size Class >>>> | | | | | | | | rge Total / cm + | | All Sizes | |
| Species | AGS | UGS | AGS | UGS | AGS | UGS | AGS | UGS | AGS | UGS | |
| Ironwood (Ostrya virginiana) | 6 | | 1 | | | | | | 7 | 0 | |
| Black Cherry (Prunus serotina) | 3 | | 1 | | | | | | 4 | 0 | |
| Sugar Maple (Acer saccharum) | 2 | | | | | | | | 2 | 0 | |
| | | | | | | | | | 0 | 0 | |
| Total Number of Trees | 11 | 0 | 2 | 0 | 0 | 0 | 0 | 0 | 13 | 0 | |

Description

| Location: | 5868 County Road 65 |
|-------------------|---------------------|
| Date: | 24-Mar-23 |
| Surveyor: | SA |
| Compartment: | P6 |
| Stations Tallied: | 100% Tally |

| | D. I. | Polewood 10-24 cm | | Sawtimber Sizes | | | | | | |
|---------------------------------------|-------|----------------------|-----|-------------------|-----|--------------------|-----|------------------|-----|-----------------|
| Tree Size Class >>>> | | | | Small 26-36 cm | | Medium 38-48 cm | | Large 50 cm + | | Total All Sizes |
| Species | AGS | UGS | AGS | UGS | AGS | UGS | AGS | UGS | AGS | UGS |
| Red Oak (Quercus rubra) | 2 | 1 | 3 | | | | | | 5 | 1 |
| Ironwood (Os <i>trya virginiana</i>) | 5 | | | 1 | | | | | 5 | 1 |
| Green Ash (Fraxinus pennsylvanica) | | 5 | | 1 | | | | | 0 | 6 |
| Sugar Maple (Acer saccharum) | 36 | 2 | 4 | | 1 | | | | 41 | 2 |
| Basswood (Tilia americana) | 1 | 2 | | | | | | | 1 | 2 |
| White Pine (Pinus strobus) | 1 | | | 1 | | | | | 1 | 1 |
| Trembling Aspen (Populus tremuloides) | 6 | 11 | 6 | | 6 | | | | 18 | 11 |
| Black Cherry (Prunus serotina) | | 3 | | | | | | 1 | 0 | 4 |
| Yellow Birch (Betula alleghaniensis) | 2 | | | | | | | | 2 | 0 |
| Pin Cherry (Prunus pensylvanica) | 3 | 5 | | | | | | | 3 | 5 |
| White Birch (Betula papyrifera) | 5 | 1 | | | | | | | 5 | 1 |
| Manitoba Maple (Acer negundo) | 3 | 3 | | | | | | | 3 | 3 |
| Black Locust (Robiniana pseudoacacia) | 4 | | 1 | | 1 | | | | 6 | 0 |
| | | | | | | | | | 0 | 0 |
| | | | | | | | | | 0 | 0 |
| Total Number of Trees | 68 | 33 | 14 | 3 | 8 | 0 | 0 | 1 | 90 | 37 |

Description

| Location: | 5868 County Road 65 |
|-------------------|---------------------|
| Date: | 24-Mar-23 |
| Surveyor: | SA |
| Compartment: | P7 |
| Stations Tallied: | 4 |

10m radius fixed area plots

| | Dala | | | | Total All Sizes | | | | | |
|---|----------------------|-----|-------------------|-----|-----------------|-----|--------------------|-----|------------------|-----|
| Tree Size Class >>>> | Polewood 10-24 cm | | Small 26-36 cm | | | | Medium 38-48 cm | | Large 50 cm + | |
| Species | AGS | UGS | AGS | UGS | AGS | UGS | AGS | UGS | AGS | UGS |
| Eastern Hemlock (<i>Tsuga canadensis</i>) | 13 | 1 | 3 | | 2 | 1 | 1 | | 19 | 2 |
| Sugar Maple (Acer saccharum) | 11 | 2 | 5 | 1 | 4 | 1 | 1 | | 21 | 4 |
| Yellow Birch (Betula alleghaniensis) | 1 | | | | | | | | 1 | 0 |
| American Beech (Fagus grandifolia) | | 3 | | 1 | | | | | 0 | 4 |
| White Ash (Fraxinus americana) | | | | | | | | | 0 | 0 |
| Ironwood (Ostrya virginiana) | | 2 | | | | | | | 0 | 2 |
| Black Cherry (Prunus serotina) | 3 | | | | | | 1 | | 4 | 0 |
| Trembling Aspen (Populus tremuloides) | | 1 | | | 1 | | | | 1 | 1 |
| White Pine (Pinus strobus) | | | | | 2 | | 1 | | 3 | 0 |
| Basswood (Tilia americana) | 2 | | 1 | | | | | | 3 | 0 |
| Red Oak (Quercus rubra) | | | | | 1 | | | | 1 | 0 |
| White Birch (Betula papyrifera) | 3 | 2 | | | | | | | 3 | 2 |
| | | | | | | | | | 0 | 0 |
| | | | | | | | | | 0 | 0 |
| Total Number of Trees | 33 | 11 | 9 | 2 | 10 | 2 | 4 | 0 | 56 | 15 |

Description:

Regeneration (<10cm DBH) within unit including Hemlock, Sugar Maple, Beech, Ironwood, Black Cheery, White Ash, Basswood,

| Location: | 5868 County Road 65 |
|-------------------|---------------------|
| Date: | 24-Mar-23 |
| Surveyor: | SA |
| Compartment: | P8 |
| Stations Tallied: | 3 |

3.99m radius fixed area plots

| Tree Size Class >>>> | Polewood 10-24 cm | | Sawtimber Sizes | | | | | | | | |
|---------------------------------------|----------------------|-----|-------------------|-----|--------------------|-----|------------------|-----|-----------------|-----|--|
| | | | Small 26-36 cm | | Medium 38-48 cm | | Large 50 cm + | | Total All Sizes | | |
| Species | AGS | UGS | AGS | UGS | AGS | UGS | AGS | UGS | AGS | UGS | |
| Trembling Aspen (Populus tremuloides) | 3 | | 1 | | | | | | 4 | 0 | |
| Ironwood (Ostrya virginiana) | 1 | | | | | | | | 1 | 0 | |
| Sugar Maple (Acer saccharum) | 1 | | | | | | | | 1 | 0 | |
| White Birch (Betula papyrifera) | 7 | | | | | | | | 7 | 0 | |
| White Elm (Ulmus americana) | 1 | | | | | | | | 1 | 0 | |
| Yellow Birch (Betula alleghaniensis) | 1 | | | | | | | | 1 | 0 | |
| | | | | | | | | | 0 | 0 | |
| Total Number of Trees | 14 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 15 | 0 | |

Description:

Dense amounts of regen-sized trees (<10cm DBH) including Trembling Aspen, Green Ash, White Birch, Black Cherry, and Sugar Maple

| Location: | 5868 County Road 65 | | | | | | |
|-------------------|---------------------|--|--|--|--|--|--|
| Date: | 24-Mar-23 | | | | | | |
| Surveyor: | SA | | | | | | |
| Compartment: | P10 | | | | | | |
| Stations Tallied: | 100% Tally | | | | | | |

| Tree Size Class >>>> Species | Data | Polewood 10-24 cm | | Sawtimber Sizes | | | | | | |
|--------------------------------------|------|----------------------|-----|-------------------|-----|--------------------|-----|------------------|-----|-----------------|
| | > | | | Small 26-36 cm | | Medium 38-48 cm | | Large 50 cm + | | Total All Sizes |
| | AGS | UGS | AGS | UGS | AGS | UGS | AGS | UGS | AGS | UGS |
| White Birch (Betula papyrifera) | 22 | | 5 | | | | | | 27 | 0 |
| Basswood (Tilia americana) | 7 | | | | | | | | 7 | 0 |
| Yellow Birch (Betula alleghaniensis) | 1 | | 1 | | | | | | 2 | 0 |
| Sugar Maple (Acer saccharum) | 13 | 1 | | | | | | | 13 | 1 |
| Silver Maple (Acer saccharinum) | 1 | | | | | | | | 1 | 0 |
| Cherry species (Prunus sp.) | 6 | | | | | | | | 6 | 0 |
| Ironwood (Ostrya virgiana) | 2 | | | | | | | | 2 | 0 |
| American Beech (Fagus grandifolia) | 1 | | | | | | | | 1 | 0 |
| Blue Beech (Carpinus caroliniana) | 2 | | | | | | | | 2 | 0 |
| Black Cherry (Prunus serotina) | 1 | | | | | | | | 1 | 0 |
| | | | | | | | | | 0 | 0 |
| | | | | | | | | | 0 | 0 |
| Total Number of Trees | 56 | 1 | 6 | 0 | 0 | 0 | 0 | 0 | 62 | 1 |

Description

Sugar Maple, Beech, and White Birch regen

| Location: | 5868 County Road 65 | | | | | | |
|-------------------|---------------------|--|--|--|--|--|--|
| Date: | 24-Mar-23 | | | | | | |
| Surveyor: | SA | | | | | | |
| Compartment: | P11 | | | | | | |
| Stations Tallied: | 100% Tally | | | | | | |

| Tree Size Class >>>> | Dala | Polewood 10-24 cm | | Sawtimber Sizes | | | | | | |
|---------------------------------------|------|----------------------|-----|-------------------|-----|--------------------|-----|------------------|-----|-----------------|
| | >> | | | Small 26-36 cm | | Medium 38-48 cm | | Large 50 cm + | | Total All Sizes |
| Species | AGS | UGS | AGS | UGS | AGS | UGS | AGS | UGS | AGS | UGS |
| Basswood (Tilia americana) | | 1 | | | | | | | 0 | 1 |
| Red Oak (Quercus rubra) | | 1 | | | | | | | 0 | 1 |
| Sugar Maple (Acer saccharum) | 12 | 1 | | | | | | | 12 | 1 |
| White Birch (Betula papyrifera) | 7 | 3 | | | | | | | 7 | 3 |
| Green Ash (Fraxinus pennsylvanica) | | 1 | | | | | | | 0 | 1 |
| Trembling Aspen (Populus tremuloides) | 5 | 5 | 1 | | | | | | 6 | 5 |
| Yellow Birch (Betula alleghaniensis) | 20 | 1 | 1 | | | | | | 21 | 1 |
| | | | | | | | | | 0 | 0 |
| | | | | | | | | | 0 | 0 |
| Total Number of Trees | 44 | 13 | 2 | 0 | 0 | 0 | 0 | 0 | 46 | 13 |

Description